



Snow analysis for NWP at Meteo France and recent developments for land surface analysis

Camille Birman, Jean-François Mahfouf...

SURFEX User Workshop, 18-20 March 2019, Toulouse, France

Outline


- Current land surface assimilation system
- Snow analysis:
 - AROME model
 - ARPEGE model
 - First steps toward the use of satellite observations
- Future plans for the assimilation system over land and preliminary steps towards coupled assimilation: Diagnostics using ARPEGE EDA for surface analysis
- Conclusions and future work

Land surface assimilation system

- Météo France global model ARPEGE and limited area model AROME are coupled to SURFEX.
- The 4 tiles have their own prognostic variables (and **analysed variables**):
 - Nature: ISBA-3L (3 layers) for NWP (Noilhan and Mahfouf, 1996; Boone et al., 1999), prognostic variables in the three superficial layers (liquid and solid fractions for soil water content, SWE for snow on the ground) → : T_s , T_2 , T_3 , w_g , w_2 , w_3 .
 - Town: TEB (Masson, 2000) → T_{roof} , T_{wall} , T_{road}
 - Lake → LST (FLake)
 - Sea → SST (CMO 1D)
- We use screen level observations of T_{2m} and RH_{2m} to compute gridded analysed fields using 2D Optimal Interpolation (OI) (CANARI)
- 1D OI scheme using the increments of T_{2m} and RH_{2m} (Giard and Bazile, 2000) to obtain increments of soil temperature and moisture T_s , T_2 , w_g , w_2 .

Snow analysis

- Snow analysis over plains: necessary to correct for insufficient snow melt in the model
 - Case study February 2018 and January-February 2019




Snow depth observations over the AROME-France domain


- Observations extracted over France
- Snow analysis performed using CANARI 2D OI
- Transfer of snow increments into SURFEX
- Prognostic variable: SWE → use of model density to transform snow depth observations into snow water equivalent for the assimilation
- Tuning of observation and background errors and length scale

Snow analysis: case study on 12th February 2018, 12:00


Background




Analysis



Reference



Analysis
Increment
($\sigma_0 = 5 \text{ kg/m}^2$)




(kg/m^2)




Snow analysis: case study on 12th February 2018, 12:00


Background





Analysis



Reference




Analysis
Increment
($\sigma_0 = 2 \text{ kg/m}^2$)



(kg/m^2)




Snow analysis: bias and MSE on February 2018 with respect to Synop stations



Snow analysis: use of satellite observations

- Daily snow product H10 from H-SAF (<http://hsaf.meteoam.it/snow.php>)
 - Resolution 0.05° over Europe, computed from SEVIRI multichannel analysis




H-SAF H10 snow product on March 9, 2010

- The snow mask (values 0 or 1) is converted into values that can be ingested in the assimilation :
 - 0 kg/m^2 (no snow) or 5 kg/m^2 (snow on the ground)
- Tuning of observation errors and length scale:
 - $\sigma_o = 8 \text{ kg/m}^2$ (H-SAF) instead of 5 kg/m^2 (Synop)
 - $d = 5 \text{ km}$ (H-SAF) instead of 50 km (Synop)


SUW, Toulouse, 18th March 2019

Snow analysis: case study on 12th February 2018, 12:00



Background




Analysis (Synop only)



Analysis (Synop + H-SAF)



Analysis Increment




Synop only

(kg/m²)
Synop + H-SAF



Snow analysis in ARPEGE

- Synop stations: heterogeneous coverage over the globe




- Same principle as in AROME: 2D-OI with a structure function accounting for the distance and the difference of elevations between observation and grid point
- Tuning of observation errors and length scale:
 - $\sigma_o = 5 \text{ kg/m}^2$, $d = 100 \text{ km}$


Snow analysis: case study on 4th February 2018, 00:00

Background

Analysis




Analysis Increment (kg/m²)



Snow analysis in ARPEGE: use of satellite data

- IMS NOAA-NESDIS snow product over Northern Hemisphere: derived from AVHRR, AMSU, GOES/Imager, Himawari (AHI), Meteosat (SEVIRI)
- Daily product, 4 km resolution
- The snow mask (values 0 or 1) is converted into values that can be ingested in the assimilation :
 - 0 kg/m² (no snow) or 5 kg/m² (snow on the ground)
- Thinning of observations: ~1 observation every 30 km is kept
- Tuning of observation errors and length scale:
 - $\sigma_o = 8$ kg/m² (IMS) instead of 5 kg/m² (Synop)
 - $d = 10$ km (IMS) instead of 100 km (Synop)




IMS snow and ice product over Northern Hemisphere on March 11, 2019


Snow analysis: case study on 20th November 2018, 00:00

Analysis


Synop only



Synop + IMS





Synop only



Analysis increment (kg/m²)

Synop + IMS



Diagnostics using ARPEGE EDA for surface analysis

- Optimal interpolation coefficients:

- Covariances between the forecast errors of T_{2m} and RH_{2m} and the perturbed soil moisture values w_g and w_2 were obtained from a set of 100 single column model integrations where the initial soil moisture content was perturbed.

$$w^a - w^b = \alpha(T^o - T^b) + \beta(RH^o - RH^b)$$

$$\alpha = \frac{\sigma_w}{\Phi \sigma_T} \left\{ \left[1 + \left(\frac{\sigma_{RH}^o}{\sigma_{RH}^b} \right)^2 \right] \rho_{T,w} - \rho_{T,RH} \rho_{RH,w} \right\}$$

$$\beta = \frac{\sigma_w}{\Phi \sigma_{RH}} \left\{ \left[1 + \left(\frac{\sigma_T^o}{\sigma_T^b} \right)^2 \right] \rho_{RH,w} - \rho_{T,RH} \rho_{T,w} \right\}$$

$$\Phi = \left[1 + \left(\frac{\sigma_T^o}{\sigma_T^b} \right)^2 \right] \left[1 + \left(\frac{\sigma_{RH}^o}{\sigma_{RH}^b} \right)^2 \right] - \rho_{T,RH}^2$$


- Kalman filter approach:

$$\Delta \mathbf{x} = \mathbf{B} \mathbf{H}^T (\mathbf{H} \mathbf{B} \mathbf{H}^T + \mathbf{R})^{-1} \Delta \mathbf{y}$$


- Use of EDA (AEARP, 25 members) to compute standard deviations and covariances between surface variables and observed variables.

$$\Delta \mathbf{x} = \text{cov}(\mathbf{x}^b, \mathbf{y}^b) \left[(\text{cov}(\mathbf{y}^b, \mathbf{y}^b) + \text{cov}(\mathbf{y}^o, \mathbf{y}^o)) \right]^{-1} \Delta \mathbf{y}$$


Diagnostics using ARPEGE EDA for surface analysis



Surface fields at **low resolution** from EDA (25 members)



Surface fields at **high resolution** (25 members)



Correlations and standard deviations at **high resolution**

Soil analysis


Diagnostics using ARPEGE EDA for surface analysis

- Cycling of analysed surface fields computed from the EDA over assimilation cycles: increments in the 2nd layer of the soil at first analysis time (20180102H00)


Ol increments

TG2

EDA increments



WG2



Conclusions and future work

- Snow analysis in AROME and ARPEGE models with a 2D Optimal Interpolation, using Synop observations
- Improvement of the land surface data assimilation system: ensemble methods (EnKF, particle filter...) and use of atmospheric ensembles produced by ensemble data assimilation systems (AEARP → global model ARPEGE, AEARO → limited area model AROME)
- Assimilation of satellite products, in particular for surface temperature (PhD Zied Sassi) and for snow (Nesdis-IMS snow cover product for ARPEGE, H-SAF products for AROME)
- Future plans for the assimilation of soil moisture products (soil moisture product from ASCAT, and/or from L-band sensors SMOS/SMAP), and albedos (LSA-SAF products)

Land surface assimilation system

- Météo France global model ARPEGE and limited area model AROME are coupled to the surface modelling platform SURFEX, which represents the exchanges between the atmosphere and the surface.
- Each grid box is divided into 4 tiles for nature, sea, lake and town fractions. The 4 tiles receive the same atmospheric forcing and the surface fluxes computed on each tile are averaged over the grid box.
- The 4 tiles have their own prognostic variables (and **analysed variables**):
 - Nature: ISBA-3L (3 layers) for NWP (Noilhan and Mahfouf, 1996; Boone et al., 1999), prognostic variables in the three superficial layers (liquid and solid fractions for soil water content, SWE for snow on the ground) → : T_s , T_2 , T_3 , w_g , w_2 , w_3 .
 - Town: TEB (Masson, 2000) → T_{roof} , T_{wall} , T_{road}
 - Lake → LST (FLake)
 - Sea → SST (CMO 1D)

Land surface assimilation system






- We use screen level observations of T_{2m} and RH_{2m} to compute gridded analysed fields using 2D Optimal Interpolation (OI) (CANARI)
- 1D OI scheme using the increments of T_{2m} and RH_{2m} (Giard and Bazile, 2000):

$$\bullet \Delta T_s = \Delta T_{2m}$$

$$\bullet \Delta T_2 = \Delta T_{2m} / 2\pi$$


$$\bullet \Delta w_g = \alpha_1 \Delta T_{2m} + \beta_1 \Delta RH_{2m}$$

$$\bullet \Delta w_2 = \alpha_2 \Delta T_{2m} + \beta_2 \Delta RH_{2m}$$




Snow analysis: case study on 22th January 2019, 12:00


Background




Analysis



Reference




Analysis
Increment
($\sigma_0 = 5 \text{ kg/m}^2$)



(kg/m²)




Snow analysis: bias and MSE on February 2018 with respect to Synop stations (9 simulations)



Bias (dashed line) and Mean Square Error (solid line) of 2 metre temperature and 2 metre relative humidity forecasts as a function of the forecast range, with (blue) and without (red) snow analysis


Diagnostics using ARPEGE EDA for surface analysis

- Diagnostics on a summer period (August 2017): Correlations between w_2 and T_{2m} averaged over the period. Coll. with ECMWF (Patricia de Rosnay)



Diagnostics using ARPEGE EDA for surface analysis

- Analysis increments for soil variables and comparison with OI increments (operationnal T_{2m} and RH_{2m} increments)



Conclusions and future work

- Analysis of surface parameters: snow depth, precipitation (2D optimal interpolation), ...(LAI); improvement of analysis techniques, in particular in areas of complex orography (MESCAN)
- Improvement of the land surface data assimilation system: ensemble methods (EnKF, particle filter...) and use of atmospheric ensembles produced by ensemble data assimilation systems (AEARP → global model ARPEGE, AEARO → limited area model AROME)
- Assimilation of satellite products, in particular for surface temperature (PhD Zied Sassi), for soil moisture (soil moisture product from ASCAT, and/or from L-band sensors SMOS/SMAP), for snow (snow cover products Nesdis-IMS/H-SAF product/Modis?) and for albedos (products from LSA-SAF)