



---

## Erratum to ‘Seasonal surface urban energy balance and wintertime stability simulated using three land-surface models in the high-latitude city Helsinki’

P. Karsisto,<sup>a,b</sup> C. Fortelius,<sup>b</sup> M. Demuzere,<sup>c</sup> C. S. B. Grimmond,<sup>d</sup> K. W. Oleson,<sup>e</sup> R. Kouznetsov,<sup>b,f</sup>  
V. Masson<sup>g</sup> and L. Järvi<sup>a\*</sup>

<sup>a</sup>*Department of Physics, University of Helsinki, Finland*

<sup>b</sup>*Finnish Meteorological Institute, Helsinki, Finland*

<sup>c</sup>*Department of Earth and Environmental Sciences, KU Leuven, Belgium*

<sup>d</sup>*Department of Meteorology, University of Reading, UK*

<sup>e</sup>*National Center for Atmospheric Research, Boulder, CO, USA*

<sup>f</sup>*Obukhov Institute for Atmospheric Physics, Moscow, Russia*

<sup>g</sup>*CNRM–GAME, Météo-France, CNRS, Toulouse and Grenoble, France*

\*Correspondence to: L. Järvi, Department of Physics, P.O. Box 00014, University of Helsinki, Finland.

E-mail: leena.jarvi@helsinki.fi

---

The above article was originally published on Early View on 9 October 2015, and subsequently in volume 142 (issue 694): 401–417; DOI:10.1002/qj.2659.

In section 5.3 *Sensible heat flux*, the numbers of RMSE are wrong in the second paragraph. The text should remain the same but the numbers in brackets should be as follows:

CLM simulates the wintertime  $Q_H$  best at Kumpula (RMSE = 27 W m<sup>-2</sup>), but overestimates  $Q_H$  at Torn

(RMSE = 73 W m<sup>-2</sup>) particularly at night. Although SUEWS is able to predict the difference between the two sites, it underestimates nocturnal  $Q_H$  at Torn and overestimates daytime  $Q_H$  at Kumpula (RMSE of 46 and 32 W m<sup>-2</sup> respectively). SURFEX underestimates  $Q_H$  at both Torn and Kumpula (RMSE = 60 and 33 W m<sup>-2</sup>, respectively) and is not able to predict the difference between the two sites in wintertime.